



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2008

LRRK2 mutations are not frequent in Swiss patients with Parkinson's disease

Burgunder, J M ; Tan, E K ; Wirthmüller, U ; Jung, H H

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-10053>

Journal Article

Accepted Version

Originally published at:

Burgunder, J M; Tan, E K; Wirthmüller, U; Jung, H H (2008). LRRK2 mutations are not frequent in Swiss patients with Parkinson's disease. *Schweizer Archiv für Neurologie und Psychiatrie*, 159(7):412-418.

LRRK2 mutations are not frequent in Swiss patients with Parkinson's disease

J.-M. Burgunder^{a, b}, E. K. Tan^c, U. Wirthmüller^d, H. H. Jung^e

^a Department of Neurology, University of Berne

^b Division of Neurology, Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore (SGP)

^c National Neuroscience Institute, SingHealth, Singapore General Hospital (SGP)

^d Institute of Immunology, University of Berne

^e Department of Neurology, University Hospital Zurich

*To whom correspondence should be addressed: Prof. Dr. med. Jean-Marc Burgunder, Neurologie Praxis, Steinerstrasse 45, 3006 Bern

Summary

A wealth of recent data has shed light on the genetics of Parkinson's disease. Mutations of specific genes have been discovered in a number of autosomal dominant and recessive forms of the disease, and polymorphisms in the same or other genes found to be associated with its occurrence in sporadic forms. Mutations in the gene encoding leucine-rich repeat kinase 2 (LRRK2) cause an autosomal dominant form of Parkinson's disease (PARK8) and have been found in patients with Parkinson's disease with a highly variable frequency in different populations in the world. In this study we have examined the most common mutations so far described in this large gene: p.R1441C (together with p.R1441G and p.R1441H), p.Y1699C and p.G2019S. Seventy-three patients from Switzerland with Parkinson's disease diagnosed according to Brain Bank criteria with the exception of positive family history were included in this study. The three exons, in which the above mutations had been found, were sequenced after amplification. There were no sequence differences compared with the published wild-type sequences in all patients. These data point at the low incidence of LRRK2 mutation in the Swiss (regions of Berne and Zurich) populations, similar to findings in other central European populations, but at great variance to Mediterranean populations.

Keywords: Parkinson; gene; LRRK2

Introduction

Parkinson's disease is a frequent disorder, 1% of people older than 60 years in Western countries are estimated to be affected, and the average age at onset is 60 [1]. With an age of 85, the prevalence even increases to 4% [2]. Environmental factors, including industrial pollutants, farming, living in a rural area, drinking well water, pesticide and herbicide exposure and head trauma [3–9] have been associated with an increased risk to develop the disorder. The best-established example of an environmental factor leading to Parkinson's disease is MPTP, an oxidant present in artisanal drug preparations, which had caused an epidemic of the disorder in young patients in the eighties of the last century, and which is now commonly used to induce Parkinson's disease in animal models [10]. Other non-genetic causes for Parkinson's disease have been reviewed recently [11].

However, a positive family history has consistently been reported as a strong risk factor for the disease [3, 12], and a large family from Grisons has been reported in the early literature in the field [13]. Up to 27% of patients with the disorder report a relative with Parkinson's disease and up to 16% its presence in a parent or sibling. The risk of first degree relatives to get Parkinson's disease is up to 9.5 times greater as compared to relatives of controls [8, 12, 14, 15]. In a large twin study monozygotic twin pairs with onset in at least one person before the age of 50 had a significant higher risk of 6 times as compared to dizygotic twin pairs [16]. When the involvement of the dopaminergic function was examined using PET, the concordance in a group of 18 monozygotic twin pairs was significant at baseline, as compared to 16 dizygotic twin pairs [17]. Furthermore, the proportion became even higher in monozygotic twins during follow-up [17]. In a more recent study similarity for age at onset among sibling pairs with Parkinson's disease and the greater risk for biological relatives as compared to spouses have again supported the presence of a genetic component [18]. Risk of Parkinson's disease was high in first relatives of both early and late-onset patients, the risk for siblings was increased only for those of early onset patients [19]. Other studies have started to examine the interaction between genetic and environmental factors. For example, there is a significant increase in risk in cases with positive family history and smoking [20].

The strongest evidence for a genetic causation of Parkinson's disease has come from the observation of families with a clear-cut autosomal dominant or recessive inheritance. The investigations in these families with the performance of linkage studies have allowed the

discovery of different genes (table 1) [21, 22]; furthermore, additional loci are known, and in time more genes with mutation directly causing the disorder are expected to be discovered.

α -synuclein was the first gene with mutations causing Parkinson's disease [23] and a p.A53T was found in one large family of Italian origin and three smaller pedigrees from Greece [23]. Later a p.A30P [24] and a p.E46K [25] mutation were found in other families of German or Spanish origin, respectively. Duplication or triplication of the region containing the gene, or of individual exons was found in additional families [26], with dementia in addition to Parkinsonian features, and a more severe phenotype in patients with triplication [27], pointing at a gene dose effect with more prominent α -synuclein accumulation within neurons and subsequent loss of dopamine function. The p.A53T mutation was found in additional families of Greek and Italian origin, most probably related to the first one, however, a number of other studies could not find additional mutations of this gene [28–31]. Even considering the very low number of cases with α -synuclein mutation, the discovery of this protein has greatly advanced our knowledge of the molecular pathophysiology of idiopathic Parkinson's disease, Lewy body disease and other neurodegenerative disorders. This protein is also involved in cases without mutations, leading to the concept of synucleinopathies [32], including the definition of potential molecular therapeutic targets [33].

Mutations in the gene coding for parkin [34] were found later, first in Japanese patients with autosomal recessive early onset parkinsonism. A large number of additional mutations were found in many other populations studied, including more than 50 point mutations, as well as exonic rearrangements, duplications and deletions, accounting for half of familial cases and 70% of sporadic cases with age of onset younger than 20 years [35–37]. The search for parkin mutations can be considered in young patients with the disorder, or in those with an autosomal recessive inheritance. Following the discovery of missense and deletion mutations in families with recessive Parkinson's disease [38], large-scales studies have examined the DJ-1 gene in other patients but only rare additional mutations have been found [39–43], suggesting that only 1% of patients with early onset Parkinson's disease may have a DJ-1 mutation.

Two mutations were found in a novel gene PINK1 [44] in families with autosomal recessive juvenile-onset Parkinson's disease. Later additional mutations, including frameshift, missense and truncating mutations were indentified [45–48], with a relative amount of 1–7% of patients with Parkinson's disease of Caucasian origin [43], suggesting that PINK1 and parkin

may be the most common genes in autosomal recessive and possibly also sporadic young-onset Parkinson's disease.

A p.I93M substitution found in exon 4 of ubiquitin carboxy-terminal hydrolase L1 (UCHL1) has been reported in a German family [49]. This has led to the examination of the role of this enzyme in Parkinson's disease, but, no further mutation has been found in numerous studies.

Further loci have been mapped in additional families (table 1), and the discovery of the respective genes is expected to further advance our understanding of the molecular pathophysiology of the disease. It is of note that mutations in other genes, involved in neurodegenerative disorders other than Parkinson's disease, may sometimes have presentations similar to this disorder. This includes frontotemporal dementia, some of the spinocerebellar ataxias, dystonia-parkinson syndrome and mitochondrial disorders.

Finally, PARK8 has been found on chromosome 12p11.2–q13 in a Japanese family with autosomal dominant Parkinson's disease with otherwise typical presentation similar to idiopathic Parkinson's disease, including age at onset, laterality and good response to L-dopa. The overwhelming majority of cases have typical Lewy body pathology, but absence of Lewy bodies in rare cases [50, 51]. This locus has been confirmed in families from other ethnic origins [52] and shortly thereafter, several mutations in a novel gene encoding leucine-rich repeat kinase 2 (LRRK2) cloned within this locus have been reported [52, 53]. Mutations in the LRRK2 account to about 13% of cases with autosomal dominant Parkinson's disease [28], they are also found in a substantial number of sporadic cases [54]. As an example of a local study we therefore examined the most commonly found LRRK2 mutations in a cohort of patients with Parkinson's disease from Switzerland.

Material and Methods

Seventy-three consecutive patients with Parkinson's disease were included. They had been examined at the Department of Neurology, University Hospital, Zurich (H.H.J.) or at the Neurology practice in downtown Berne (J.-M. B.) and were all of Caucasian origin. The study has been approved by the respective Ethics Committees of the Cantons of Zurich and Berne. The diagnosis had been made according to Brain Bank criteria, with the exception of positive family

history as an exclusion criterion for idiopathic Parkinson's disease, and all patients responded well to treatment with L-dopa or dopamine agonists.

Blood was withdrawn and DNA extracted according to standard procedures using QIAGEN PCR purification kit. Following primer pairs were used: for exon 31, mutations R1441C, R1441G and R1441H: 5'-CTTCTGAAGTCTGCTAGTTTCT-3'(forward) and 5'-CGTTGGCACACATTTGGATG-3'(reverse); for exon 35, mutation Y1699C: 5'-GCTCAACAAGGTTGGGTGTT-3'(forward) and 5'-ATGCTTTCAGGGAGAGGTAAG-3'(reverse); for exon 41, mutation G2019S: 5'-TTTTGATGCTTGACATAGTGGAC-3'(forward) and 5'-CACATCTGAGGTCAGTGGTTATC-3'(reverse). Direct sequencing was performed after optimised PCR amplification using Big Dye Terminator ver.3.1 cycle (Applied Biosystems) on an ABI3100 Genetic Analyser and analysed with DNA Sequencing Analysis (ver.3.7) software (Applied Biosystems).

Results

The age of the patients varied from 49 to 81 years (mean 70), 43% were women and 17% had a positive family history. No mutation was detected in any of the three exons studied in any of the patients. No polymorphisms were found either.

Discussion

Our exploration in a preliminary small set of patients with idiopathic and familial Parkinson's disease from Switzerland did not demonstrate the presence of any of the LRRK2 mutations most commonly found in similar or larger sets of patients from other countries. Due to the small size of the sample, we cannot exclude the presence of these mutations in PD patients from Swiss origin, but based on our data we can conclude that the frequency will most probably be in the lower range found in different populations. A high frequency of the p.G2019S mutation in sporadic Parkinson's disease has been found in North African Arabs (20%) [55] and Ashkenazi Jews (13%) [56], where- as the same mutation was absent in Chinese PD patients [57]. About 7% of PD patients from cohorts of similar size as ours in the Iberic peninsula carried the mutation [58, 59], and a low prevalence of this mutation from 0 to 3% was documented in other European

populations, like Poland [60], Norway [61] and Germany [28]. The absence of this mutation in our set of patients is in line with the low frequency of the mutation in North and Middle European populations. As the penetrance of the p.G2019S mutation increases with age [54, 62] and the age of patients in our cohort is in the higher range, the absence of this mutation was not due to a selection bias of younger patients. Several mutations at position 1441 have been described earlier in different populations. The p.R1441H mutation has been found in rare families of different origins [59, 63–65]. In contrast, the p.R1441G mutation is quite frequent in PD patients from the Basque region [66].

LRRK2 codes for a protein with 2527 amino acids [52], which has a molecular mass of approximately 250–280 kDA [67, 68]. The gene is selectively expressed at high levels in the mouse [69], rat and human striatum [70], which is in contrast with the more widespread expression of other genes mutated in familial forms of Parkinson's disease. The presence of specific domains, including ANK, LRR, Roc, COR, Kinase or MAPKKK, and WD40, suggests a role in signalling [52, 71]. The soluble protein is located in the cytoplasm [72], consistent with the absence of specific targeting signal, as demonstrated after HA-tagging in cell culture and immunohistochemistry in human brain sections [68]. Subcellular fractionation suggests a location close to mitochondria, microtubules, Golgi and endoplasmic reticulum [73]. Interactions of LRRK2 have been demonstrated with parkin, but not with DJ-1, α -synuclein or tau [72]. Mutations of the protein do not consequently give rise to their accumulation [68], which is in contrast with the clear-cut pathophysiological mechanism of α -synuclein accumulation in hereditary and sporadic Parkinson's disease. Functional analysis of one mutation, p.I2020T, located at the beginning of the kinase domain activation loop, led to the description of a significantly higher kinase activity, pointing at a gain-of-function for this particular mutation [73]. The p.R1441C, p.Y1699C and p.G2019S mutations, which we have studied in the present report, lead to neurodegeneration when expressed in specific cells in culture [72]. The p.R1441C mutation is localised within the GTPase domain, and the p.G2019S within the mixed kinase-like domain. Both mutations also lead to an increase in enzymatic activity, with increased substrate as well as auto-phosphorylation [67]. However, analysis of further mutations of this large protein with several domains will be needed to draw general conclusions about its physiological role and the biochemical changes induced by its mutations.

Overall, there is a wide range of clinical presentations and neuropathological findings between the different mutations, and no clear-cut molecular–clinical correlation can be drawn so

far. Although LRRK2 mutations may play a role in predisposing neurons to toxic effects, the exact nature of such epigenetic and environmental factors is not yet known. The high frequency of mutations in certain populations, both in familial and sporadic cases, indicates that this protein may play a key role in the pathogenesis of Parkinson's disease. As the other proteins involved in the pathogenetic cascade in familial Parkinsonian syndromes, LRRK2 has the potential to deliver future drug targets for disease-modifying therapies. In order to advance in that direction, further data for clinical and molecular correlation are needed, as well as further studies in the biology of this protein.

Table

Selected genes and loci associated with Parkinson's disease and parkinsonism

locus	chromosomal position	name	MIM	mode of inheritance	changes	reference
PARK01/04	4q21–23	SNCA	168601	AD	p.A30P, p.E46K, p.A53T, genomic duplication, triplications	[23]
PARK02	6q25.2–27	Parkin	600116	AR	missense, truncating mutations, exon deletions, duplications, triplications	[34]
PARK03	2p13	unknown	602404	AD	–	[74]
PARK05	4p14–15	UCH L1	191342	AD	one mutation	[49]
PARK06	1p35–p36	PINK1	605909	AR	missense mutations, exon deletions	[44]
PARK07	1p36	DJ1	606324	AR	missense mutations, exon deletions	[38]
PARK08	12p11.2–q13.1	LRRK2	607060	AD	missense mutations	[53]
PARK09	1p	ATP13A2	600161	AR	missense mutations	[75]
PARK10	1p32	unknown	606852	AD	–	[76]
PARK11	2q36–q36	GIGYF2/ TNRC15	607688	?	missense mutations	[77]
PARK12	Xq21–q25	unknown	300557	?	–	[78]
PARK13	2p12	Omi/HTRA2	610297	AR	missense mutation	[79]

References.

- 1 Mayeux R, Marder K, Cote LJ, Denaro J, Hemenegildo N, Mejia H, et al. The frequency of idiopathic Parkinson's disease by age, ethnic group, and sex in northern Manhattan, 1988–1993. *Am J Epidemiol.* 1995;142:820–7.
- 2 de Rijk MC, Launer LJ, Berger K, Breteler MM, Dartigues JF, Baldereschi M, et al. Prevalence of Parkinson's disease in Europe: a collaborative study of population-based cohorts. Neurologic Diseases in the Elderly Research Group. *Neurology.* 2000;54(11 Suppl 5):S21–3.
- 3 Semchuk KM, Love EJ, Lee RG. Parkinson's disease and exposure to agricultural work and pesticide chemicals. *Neurology.* 1992;42:1328–35.
- 4 Golbe LI, Farrell TM, Davis PH. Follow-up study of earlylife protective and risk factors in Parkinson's disease. *Mov Disord.* 1990;5:66–70.
- 5 Semchuk KM, Love EJ, Lee RG. Parkinson's disease and exposure to rural environmental factors: a population-based case-control study. *Can J Neurol Sci.* 1991;18:279–86.
- 6 Rybicki BA, Johnson CC, Uman J, Gorell JM. Parkinson's disease mortality and the industrial use of heavy metals in Michigan. *Mov Disord.* 1993;8:87–92.
- 7 Seidler A, Hellenbrand W, Robra BP, Vieregge P, Nischan P, Joerg J, et al. Possible environmental, occupational, and other etiologic factors for Parkinson's disease: a case-control study in Germany. *Neurology.* 1996;46:1275–84.
- 8 Semchuk KM, Love EJ, Lee RG. Parkinson's disease: a test of the multifactorial etiologic hypothesis. *Neurology.* 1993;43:1173–80.
- 9 Factor SA, Weiner WJ. Prior history of head trauma in Parkinson's disease. *Mov Disord.* 1991;6:225–9.
- 10 Smeyne RJ, Jackson-Lewis V. The MPTP model of Parkinson's disease. *Brain Res Mol Brain Res.* 2005;134:57–66.
- 11 Chade AR, Kasten M, Tanner CM. Nongenetic causes of Parkinson's disease. *J Neural Transm Suppl.* 2006;(70):147–51.
- 12 Taylor CA, Saint-Hilaire MH, Cupples LA, Thomas CA, Burchard AE, Feldman RG, et al. Environmental, medical, and family history risk factors for Parkinson's disease: a New England-based case control study. *Am J Med Genet.* 1999;88:742–9.
- 13 Branger F. Une forme familiale de paralysie agitante dans une souche des grisons. *J Génét Hum.* 1956;5:261–70.
- 14 Marder K, Tang MX, Mejia H, Alfaro B, Cote L, Louis E, et al. Risk of Parkinson's disease among first-degree relatives: a community-based study. *Neurology.* 1996;47:155–60.
- 15 Payami H, Larsen K, Bernard S, Nutt J. Increased risk of Parkinson's disease in parents and siblings of patients. *Ann Neurol.* 1994;36:659–61.
- 16 Tanner CM, Ottman R, Goldman SM, Ellenberg J, Chan P, Mayeux R, et al. Parkinson's disease in twins: an etiologic study. *JAMA.* 1999;281:341–6.
- 17 Piccini P, Burn DJ, Ceravolo R, Maraganore D, Brooks DJ. The role of inheritance in sporadic Parkinson's disease: evidence from a longitudinal study of dopaminergic function in twins. *Ann Neurol.* 1999;45:577–82.
- 18 Maher NE, Golbe LI, Lazzarini AM, Mark MH, Currie LJ, Wooten GF, et al. epidemiologic study of 203 sibling pairs with Parkinson's disease: the GenePD study. *Neurology.* 2002;58:79–84.

- 19 Marder K, Levy G, Louis ED, Mejia-Santana H, Cote L, Andrews H, et al. Familial aggregation of early- and late onset Parkinson's disease. *Ann Neurol*. 2003;54:507–13.
- 20 Elbaz A, Manubens-Bertran JM, Baldereschi M, Breteler MM, Grigoletto F, Lopez-Pousa S, et al. Parkinson's disease, smoking, and family history. EUROPARKINSON Study Group. *J Neurol*. 2000;247:793–8.
- 21 Klein C, Schlossmacher MG. The genetics of Parkinson's disease: implications for neurological care. *Nat Clin Pract Neurol*. 2006;2:136–46.
- 22 Tan EK, Skipper LM. Pathogenic mutations in Parkinson's disease. *Hum Mutat*. 2007;28:641–53.
- 23 Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, et al. Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science*. 1997;276:2045–7.
- 24 Kruger R, Kuhn W, Muller T, Woitalla D, Graeber M, Kosel S, et al. Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat Genet*. 1998;18:106–8.
- 25 Zarranz JJ, Alegre J, Gomez-Esteban JC, Lezcano E, Ros R, Ampuero I, et al. The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia. *Ann Neurol*. 2004;55:164–73.
- 26 Nishioka K, Hayashi S, Farrer MJ, Singleton AB, Yoshino H, Imai H, et al. Clinical heterogeneity of alphasynuclein gene duplication in Parkinson's disease. *Ann Neurol*. 2006;59:298–309.
- 27 Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, et al. Alpha-synuclein locus triplication causes Parkinson's disease. *Science*. 2003;302:841.
- 28 Berg D, Schweitzer K, Leitner P, Zimprich A, Lichtner P, Belcredi P, et al. Type and frequency of mutations in the LRRK2 gene in familial and sporadic Parkinson's disease. *Brain*. 2005;128:3000–11.
- 29 Berg D, Niwar M, Maass S, Zimprich A, Moller JC, Wuellner U, et al. Alpha-synuclein and Parkinson's disease: implications from the screening of more than 1,900 patients. *Mov Disord*. 2005;20:1191–4.
- 30 Johnson J, Hague SM, Hanson M, Gibson A, Wilson KE, Evans EW, et al. SNCA multiplication is not a common cause of Parkinson's disease or dementia with Lewy bodies. *Neurology*. 2004;63:554–6.
- 31 Gispert S, Trenkwalder C, Mota-Vieira L, Kostic V, Auburger G. Failure to find alpha-synuclein gene dosage changes in 190 patients with familial Parkinson's disease. *Arch Neurol*. 2005;62:96–8.
- 32 Giorgi FS, Bandettini di Poggio A, Battaglia G, Pellegrini A, Murri L, Ruggieri S, et al. A short overview on the role of alpha-synuclein and proteasome in experimental models of Parkinson's disease. *J Neural Transm Suppl*. 2006;(70):105–9.
- 33 Lee VM, Trojanowski JQ. Mechanisms of Parkinson's disease linked to pathological alpha-synuclein: new targets for drug discovery. *Neuron*. 2006;52:33–8.
- 34 Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, et al. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature*. 1998;392:605–8.
- 35 Lucking CB, Durr A, Bonifati V, Vaughan J, De Michele G, Gasser T, et al. Association between early-onset Parkinson's disease and mutations in the parkin gene. French Parkinson's Disease Genetics Study Group. *N Engl J Med*. 2000;342:1560–7.
- 36 Periquet M, Lucking C, Vaughan J, Bonifati V, Durr A, De Michele G, et al. Origin of the mutations in the parkin gene in Europe: exon rearrangements are independent recurrent

- events, whereas point mutations may result from Founder effects. *Am J Hum Genet.* 2001;68:617–26.
- 37 Mata IF, Lockhart PJ, Farrer MJ. Parkin genetics: one model for Parkinson's disease. *Hum Mol Genet.* 2004;13 Spec No 1:R127–33.
 - 38 Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, et al. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science.* 2003;299:256–9.
 - 39 Lockhart PJ, Bounds R, Hulihan M, Kachergus J, Lincoln S, Lin CH, et al. Lack of mutations in DJ-1 in a cohort of Taiwanese ethnic Chinese with early-onset parkinsonism. *Mov Disord.* 2004;19:1065–9.
 - 40 Clark LN, Afridi S, Mejia-Santana H, Harris J, Louis ED, Cote LJ, et al. Analysis of an early-onset Parkinson's disease cohort for DJ-1 mutations. *Mov Disord.* 2004;19:796–800.
 - 41 Hering R, Petrovic S, Mietz EM, Holzmann C, Berg D, Bauer P, et al. Extended mutation analysis and association studies of Nurr1 (NR4A2) in Parkinson's disease. *Neurology.* 2004;62:1231–2.
 - 42 Tan EK, Tan C, Zhao Y, Yew K, Shen H, Chandran VR et al. Genetic analysis of DJ-1 in a cohort Parkinson's disease patients of different ethnicity. *Neurosci Lett.* 2004;367:109–12.
 - 43 Healy DG, Abou-Sleiman PM, Valente EM, Gilks WP, Bhatia K, Quinn N, et al. DJ-1 mutations in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2004;75:144–5.
 - 44 Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, et al. Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science.* 2004;304:1158–60.
 - 45 Tan EK, Yew K, Chua E, Puvan K, Shen H, Lee E, et al. PINK1 mutations in sporadic early-onset Parkinson's disease. *Mov Disord.* 2006;21:789–93.
 - 46 Fung HC, Chen CM, Hardy J, Singleton AB, Lee-Chen GJ, Wu YR. Analysis of the PINK1 gene in a cohort of patients with sporadic early-onset parkinsonism in Taiwan. *Neurosci Lett.* 2006;394:33–6.
 - 47 Bonifati V, Rohe CF, Breedveld GJ, Fabrizio E, De Mari M, Tassorelli C, et al. Early-onset parkinsonism associated with PINK1 mutations: frequency, genotypes, and phenotypes. *Neurology.* 2005;65:87–95.
 - 48 Albanese A, Valente EM, Romito LM, Bellacchio E, Elia AE, Dallapiccola B. The PINK1 phenotype can be indistinguishable from idiopathic Parkinson's disease. *Neurology.* 2005;64:1958–60.
 - 49 Leroy E, Boyer R, Auburger G, Leube B, Ulm G, Mezey E, et al. The ubiquitin pathway in Parkinson's disease. *Nature.* 1998;395:451–2.
 - 50 Funayama M, Hasegawa K, Kowa H, Saito M, Tsuji S, Obata F. A new locus for Parkinson's disease (PARK8) maps to chromosome 12p11.2-q13.1. *Ann Neurol.* 2002;51:296–301.
 - 51 Funayama M, Hasegawa K, Ohta E, Kawashima N, Komiyama M, Kowa H, et al. An LRRK2 mutation as a cause for the parkinsonism in the original PARK8 family. *Ann Neurol.* 2005;57:918–21.
 - 52 Zimprich A, Biskup S, Leitner P, Lichtner P, Farrer M, Lincoln S, et al. Mutations in LRRK2 cause autosomaldominant parkinsonism with pleomorphic pathology. *Neuron.* 2004;44:601–7.

- 53 Paisan-Ruiz C, Jain S, Evans EW, Gilks WP, Simon J, van der Brug M, Lopez de Munain A, et al. Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron*. 2004;44:595–600.
- 54 Kachergus J, Mata IF, Hulihan M, Taylor JP, Lincoln S, Aasly J, et al. Identification of a novel LRRK2 mutation linked to autosomal dominant parkinsonism: evidence of a common founder across European populations. *Am J Hum Genet*. 2005;76:672–80.
- 55 Lesage S, Durr A, Tazir M, Lohmann E, Leutenegger AL, Janin S, et al. LRRK2 G2019S as a cause of Parkinson's disease in North African Arabs. *N Engl J Med*. 2006;354:422–3.
- 56 Ozelius LJ, Senthil G, Saunders-Pullman R, Ohmann E, Deligtisch A, Tagliati M, et al. LRRK2 G2019S as a cause of Parkinson's disease in Ashkenazi Jews. *N Engl J Med*. 2006;354:424–5.
- 57 Tan EK, Shen H, Tan LC, Farrer M, Yew K, Chua E, et al. The G2019S LRRK2 mutation is uncommon in an Asian cohort of Parkinson's disease patients. *Neurosci Lett*. 2005;384:327–9.
- 58 Infante J, Rodriguez E, Combarros O, Mateo I, Fontalba A, Pascual J, et al. LRRK2 G2019S is a common mutation in Spanish patients with late-onset Parkinson's disease. *Neurosci Lett*. 2006;395:224–6.
- 59 Ferreira JJ, Guedes LC, Rosa MM, Coelho M, van Doeselaar M, Schweiger D, et al. High prevalence of LRRK2 mutations in familial and sporadic Parkinson's disease in Portugal. *Mov Disord*. 2007;22:1194–201.
- 60 Bialecka M, Hui S, Klodowska-Duda G, Opala G, Tan EK, Drozdik M. Analysis of LRRK2 G2019S and I2020T mutations in Parkinson's disease. *Neurosci Lett*. 2005;390:1–3.
- 61 Aasly JO, Toft M, Fernandez-Mata I, Kachergus J, Hulihan M, White LR, et al. Clinical features of LRRK2-associated Parkinson's disease in central Norway. *Ann Neurol*. 2005;57:762–5.
- 62 Clark LN, Wang Y, Karlins E, Saito L, Mejia-Santana H, Harris J, et al. Frequency of LRRK2 mutations in early and late-onset Parkinson's disease. *Neurology*. 2006;67:1786–91.
- 63 Zabetian CP, Samii A, Mosley AD, Roberts JW, Leis BC, Yearout D, et al. A clinic-based study of the LRRK2 gene in Parkinson's disease yields new mutations. *Neurology*. 2005;65:741–4.
- 64 Mata IF, Taylor JP, Kachergus J, Hulihan M, Huerta C, Lahoz C, et al. LRRK2 R1441G in Spanish patients with Parkinson's disease. *Neurosci Lett*. 2005;382:309–11.
- 65 Spanaki C, Latsoudis H, Plaitakis A. LRRK2 mutations on Crete: R1441H associated with PD evolving to PSP. *Neurology*. 2006;67:1518–9.
- 66 Paisan-Ruiz C, Saenz A, Lopez de Munain A, Marti I, Martinez Gil A, Marti-Masso JF, et al. Familial Parkinson's disease: clinical and genetic analysis of four Basque families. *Ann Neurol*. 2005;57:365–72.
- 67 West AB, Moore DJ, Choi C, Andrabi SA, Li X, Dikeman D, et al. Parkinson's disease-associated mutations in LRRK2 link enhanced GTP-binding and kinase activities to neuronal toxicity. *Hum Mol Genet*. 2007;16:223–32.
- 68 Giasson BI, Covy JP, Bonini NM, Hurtig HI, Farrer MJ, Trojanowski JQ, et al. Biochemical and pathological characterization of Lrrk2. *Ann Neurol*. 2006;59:315–22.
- 69 Melrose HL, Kent CB, Taylor JP, Dachsel JC, Hinkle KM, Lincoln SJ, et al. A comparative analysis of leucine-rich repeat kinase 2 (Lrrk2) expression in mouse brain and Lewy body disease. *Neuroscience*. 2007;147:1047–58.

- 70 Galter D, Westerlund M, Carmine A, Lindqvist E, Sydow O, Olson L. LRRK2 expression
linked to dopamine-innervated areas. *Ann Neurol*. 2006;59:714–9.
- 71 Paisan-Ruiz C, Lang AE, Kawarai T, Sato C, Salehi-Rad S, Fisman GK, et al. LRRK2
gene in Parkinson's disease: mutation analysis and case control association study.
Neurology. 2005;65:696–700.
- 72 Smith WW, Pei Z, Jiang H, Moore DJ, Liang Y, West AB, et al. Leucine-rich repeat
kinase 2 (LRRK2) interacts with parkin, and mutant LRRK2 induces neuronal
degeneration. *Proc Natl Acad Sci U S A*. 2005;102:18676–81.
- 73 Gloeckner CJ, Kinkl N, Schumacher A, Braun RJ, O'Neill E, Meitinger T, et al. The
Parkinson's disease causing LRRK2 mutation I2020T is associated with increased kinase
activity. *Hum Mol Genet*. 2006;15:223–32.
- 74 Gasser T, Muller-Myhsok B, Wszolek ZK, Oehlmann R, Calne DB, Bonifati V, et al. A
susceptibility locus for Parkinson's disease maps to chromosome 2p13. *Nat Genet*.
1998;18:262–5.
- 75 Ramirez A, Heimbach A, Grundemann J, Stiller B, Hampshire D, Cid LP, et al.
Hereditary parkinsonism with dementia is caused by mutations in ATP13A2, encoding a
lysosomal type 5 P-type ATPase. *Nat Genet*. 2006;38:1184–91.
- 76 Li YJ, Scott WK, Hedges DJ, Zhang F, Gaskell PC, Nance MA, et al. Age at onset in two
common neurodegenerative diseases is genetically controlled. *Am J Hum Genet*.
2002;70:985–93.
- 77 Lautier C, Goldwurm S, Dürr A, Giovannone B, Tsiras WG, Pezzoli G, et al. Mutations
in the GIGYF2 (TNRC15) gene at the PARK11 locus in familial Parkinson's disease. *Am
J Hum Genet*. 2008;82:822–33.
- 78 Pankratz N, Nichols WC, Uniacke SK, Halter C, Rudolph A, Shults C, et al. Genome
screen to identify susceptibility genes for Parkinson's disease in a sample without parkin
mutations. *Am J Hum Genet*. 2002;71:124–35.
- 79 Strauss KM, Martins LM, Plun-Favreau H, Marx FP, Kautzmann S, Berg D, et al. Loss of
function mutations in the gene encoding Omi/HtrA2 in Parkinson's disease. *Hum Mol
Genet*. 2005;14:2099–111.